

Transendothelial Migration of Melanoma Cells Involves N-Cadherin-mediated Adhesion and Activation of the β -Catenin Signaling Pathway

Jianfei Qi,^{*†} Ning Chen,^{*†} Junfu Wang,^{*‡} and Chi-Hung Siu^{*†}

^{*}Banting and Best Department of Medical Research and [†]Department of Biochemistry, University of Toronto, Toronto, Ontario M5G 1L6, Canada

Submitted March 7, 2005; Revised June 8, 2005; Accepted June 22, 2005

Monitoring Editor: Asma Nusrat

Cancer metastasis is a multistep process involving many types of cell-cell interactions, but little is known about the adhesive interactions and signaling events during extravasation of cancer cells. Transendothelial migration of cancer cells was investigated using an in vitro assay, in which melanoma cells were seeded on top of a monolayer of endothelial cells. Attachment of melanoma cells on the endothelium induced a twofold increase in N-cadherin expression in melanoma cells and the redistribution of N-cadherin to the heterotypic contacts. Transendothelial migration was inhibited when N-cadherin expression was repressed by antisense RNA, indicating a key role played by N-cadherin. Whereas N-cadherin and β -catenin colocalized in the contact regions between melanoma cells and endothelial cells during the initial stages of attachment, β -catenin disappeared from the heterotypic contacts during transmigration of melanoma cells. Immunolocalization and immunoprecipitation studies indicate that N-cadherin became tyrosine-phosphorylated, resulting in the dissociation of β -catenin from these contact regions. Concomitantly, an increase in the nuclear level of β -catenin occurred in melanoma cells, together with a sixfold increase in β -catenin-dependent transcription. Transendothelial migration was compromised in cells expressing a dominant-negative form of β -catenin, thus supporting a regulatory role of β -catenin signaling in this process.

INTRODUCTION

Cancer metastasis is a complex multistep process that involves the detachment of cancer cells from the primary tumor mass, intravasation, extravasation, and the establishment of new foci in a remote organ (Fidler, 2003; Pantel and Brakenhoff, 2004). All these steps involve intricate interactions between different types of cells and it is, therefore, evident that cell adhesion molecules (CAMs) play an important role in cancer metastasis. Changes in CAM profile are often associated with the disruption of normal cell-cell interactions and the establishment of new interactions, both of which are essential to metastasis formation (Christofori, 2003).

One of the least known steps in cancer metastasis is extravasation. Unlike leukocytes, only a few cancer cell types undergo rolling on the endothelial surface under flow conditions in in vitro assays (Giavazzi *et al.*, 1993; Brenner *et al.*, 1995; Aigner *et al.*, 1998). On the other hand, intravital microscopy shows that initial arrest of cancer cells occurs primarily by size restriction in the capillaries (Chambers *et al.*, 1992) and rolling has not been observed (Orr *et al.*, 2000). To investigate the mechanism of transendothelial migration, we have established an in vitro assay by depositing melanoma cells on top of a monolayer of microvascular endothelial

cells cultured on Matrigel (Sandig *et al.*, 1997b; Voura *et al.*, 1998). Adhesion among endothelial cells is mediated by several major CAMs, including VE-cadherin and PECAM-1 (Vestweber, 2002; Ilan and Madri, 2003). The attachment of melanoma cells on an endothelial monolayer has been found to induce localized dissolution of both VE-cadherin and PECAM-1 in the endothelial junction (Sandig *et al.*, 1997b; Voura *et al.*, 2000). Thus, neither VE-cadherin nor PECAM-1 appears to be involved in the transendothelial migration of melanoma cells.

Transendothelial migration is a dynamic process that involves the constant breaking and remaking of intercellular contacts and is accompanied by drastic changes in cell shape and cytoskeletal reorganization in both the tumor cell and its neighboring endothelial cells (Voura *et al.*, 1998; Brandt *et al.*, 1999). We have found that the cell adhesion molecule L1 and integrin $\alpha v \beta 3$ play a role in the formation of heterotypic contacts between melanoma cells and endothelial cells (Voura *et al.*, 2001). However, antibody and peptide inhibition studies suggest the involvement of multiple CAMs. A potential candidate is N-cadherin, because transendothelial migration can be retarded by antibodies against N-cadherin (Sandig *et al.*, 1997b).

Tumor cells from prostate, breast, and skin have been found to express high levels of N-cadherin (Islam *et al.*, 1996; Nieman *et al.*, 1999; Tomita *et al.*, 2000; Li *et al.*, 2001). In the case of melanoma progression, metastasis is accompanied by the down-regulation of E-cadherin and the up-regulation of N-cadherin expression, which facilitate the separation of melanoma cells from adjacent E-cadherin-expressing keratinocytes and the invasion of the dermal tissue through interactions with the N-cadherin-expressing fibroblasts (Li and Herlyn, 2000). Because blood vessel endothelial cells

This article was published online ahead of print in *MBC in Press* (<http://www.molbiolcell.org/cgi/doi/10.1091/mbc.E05-03-0186>) on June 29, 2005.

[†] Present address: Shandong Academy of Medical Sciences, Jinan 250062, Shandong Province, PR China.

Address correspondence to: Chi-Hung Siu (chi.hung.siu@utoronto.ca).

also express N-cadherin (Salomon *et al.*, 1992; Navarro *et al.*, 1998), it is likely that N-cadherin-dependent interactions may contribute to the adhesive events during intravasation and extravasation of melanoma cells.

N-cadherin and several members of the cadherin family mediate cell-cell adhesion via homophilic binding and the stability of cadherin-mediated cell adhesion depends on the association of β -catenin with the cadherin cytoplasmic domain. β -catenin binds α -catenin, which links the cadherin complex to the actin cytoskeleton (Takeichi, 1995; Nagafuchi, 2001). Another catenin, p120^{cas}, binds to the juxtamembrane region of the cadherin cytoplasmic domain (Anastasiadis and Reynolds, 2000).

In addition to its role in cell-cell adhesion, β -catenin is a key signal transducer in the Wnt signaling pathway, which plays an important role in many biological processes such as embryogenesis, tumorigenesis, and cancer metastasis (Smalley and Dale, 1999; Peifer and Polakis, 2000; Nelson and Nusse, 2004). Activation of Wnt signaling leads to the stabilization and translocation of β -catenin into the nucleus, where it binds the TCF/LEF family of transcription factors to induce gene transcription (Behrens *et al.*, 1996). To investigate the potential role of N-cadherin during cancer cell extravasation, immunofluorescence staining was used to examine the distribution of N-cadherin during transendothelial migration. A novel type of N-cadherin adhesion complex that lacked β -catenin was discovered in the heterotypic contact. Further studies indicate that N-cadherin became tyrosine-phosphorylated during the transmigration process, resulting in the dissociation of β -catenin, which then translocated to the nucleus of melanoma cells to activate gene transcription.

MATERIALS AND METHODS

Antibodies

Mouse mAbs against human N-cadherin (clone 32), E-cadherin, VE-cadherin, β -catenin, γ -catenin, and p120^{cas}, and anti-phospho-tyrosine monoclonal antibody (mAb; PY-20) were purchased from Transduction Laboratories (Lexington, KY). Rabbit anti-human β -catenin antibody, mouse anti-N-cadherin mAb (GC4), mouse anti-Protein-A mAb, and rabbit PAP (peroxidase-antiperoxidase) antibody were purchased from Sigma (St. Louis, MO). Mouse mAb against the dephosphorylated form of β -catenin was purchased from AG Scientific (San Diego, CA) for immunostaining of nuclear β -catenin. For immunoprecipitation studies, rabbit anti-human N-cadherin antibody (H-36) was purchased from R&D Systems (Minneapolis, MN). The Alexa488- or Alexa568-conjugated goat anti-mouse and goat anti-rabbit antibodies were purchased from Molecular Probes (Eugene, OR). Phycoerythrin (PE)-conjugated goat anti-mouse F(ab')₂ was purchased from Caltag Laboratories (Burlingame, CA).

Cell Lines

Human lung microvascular endothelial cells (HMVEC) were purchased from Clonetics (San Diego, CA) and cultured in endothelial medium (EGM-2 MV; Clonetics) supplemented with penicillin-streptomycin (100 U/ml and 100 μ g/ml, respectively). Melanoma cells were routinely cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum and penicillin-streptomycin.

cDNA Constructs

The pPUR vector that contains an N-cadherin antisense sequence was a kind gift from Dr. Keith Johnson (University of Toledo, Toledo, OH; Islam *et al.*, 1996). The original TAP-tag construct PBS1761 was obtained from Dr. Bertrand Seraphin (CNRS, France; Rigaut *et al.*, 1999). The β -catenin cDNA was obtained by RT-PCR using WM239 total RNA. The PCR product was cloned into the *Xba*I/*Kpn*I sites of the pBSK(+) vector, which was used as the PCR template to produce the truncation mutants of β -catenins (Δ N β -catenin with deletion of amino acids 1–148; Δ N+ Δ C β -catenin with deletion of amino acids 1–148 and 665–781). The PCR products were subcloned into the *Xba*I/*Kpn*I sites of the pcDNA3.1 (–) vector that contained an N-terminal TAP-tag. All the cDNA constructs were confirmed by DNA sequencing.

Cell Transfection

Plasmid DNA was transfected into WM239 melanoma cells using Lipofectamine 2000 (Invitrogen, Carlsbad, CA). The selection media were applied after 48 h of transfection (1 mg/ml G418 for the pcDNA vector; 5 μ g/ml puromycin for the pPUR vector). Stable colonies were expanded and screened by immunoblot using the PAP antibody for TAP-tag transfection and the N-cadherin mAb for N-cadherin antisense transfection.

Transendothelial Migration Assay

The in vitro transendothelial migration assay was carried out as described previously (Sandig *et al.*, 1997b; Voura *et al.*, 2001). HMVEC cells (1.5×10^5 cells, passages 5–8) in 200 μ l of endothelial medium were placed on top of a Matrigel-coated glass coverslip (12-mm diameter) and allowed to settle for 5–6 h. Coverslips were then transferred to a new 24-well plate, and incubated in 0.5 ml of endothelial medium containing 10 ng/ml TNF- α (Sigma) for 12 h to stimulate the expression of cell adhesion molecules, which in turn promote the attachment of melanoma cells (Luscinskas *et al.*, 1995; Orr *et al.*, 2000). The monolayer was washed three times and then incubated in 0.5 ml of fresh endothelial medium without TNF- α . Melanoma cells were labeled with 10 μ g/ml DiI (Molecular Probes) for 5 min at 37°C. After washing, cells were resuspended at 2.5×10^6 cells/ml in the endothelial medium. Melanoma cells (5×10^4 cells) were added to each HMVEC monolayer. Cocultures were fixed at different time points with paraformaldehyde and stained for F-actin. Quantitative analysis of transmigration was carried out using a Wild Leitz Orthoplan epifluorescence microscope (Leica Microsystems, Richmond Hill, Ontario, Canada). For each coverslip, three sets of random fields for a total of 45 fields were scored for transmigrated cells. Each set of 15 fields contained 100–200 melanoma cells and >1000 melanoma cells were routinely scored for each experimental condition.

In inhibition studies, cocultures were carried in the presence of genistein (10 μ M), calphostin C (1 μ M) or H8 (10 μ M). In antibody inhibition studies, melanoma cells were preincubated with the anti-N-cadherin function blocking antibody GC4 (1/10 dilution; Sigma) for 30 min before adding to a HMVEC monolayer. In all inhibition studies, the total number of melanoma cells associated with the endothelial monolayer was estimated to ensure that any reduction in the number of transmigrated cells was not due to a reduction in cell attachment.

Immunofluorescence Staining

Melanoma cells were labeled with Cell Tracker Orange (Molecular Probes) and then deposited on a monolayer of HMVEC. Cocultures were fixed using either 100% methanol (prechilled to –20°C) on ice for 3 min or 3.5% paraformaldehyde at room temperature for 5 min, followed by 0.1% Triton X-100 extraction for 5 min. All fixed specimens were washed and then blocked in 1% bovine serum albumin for 30 min. Incubation with the primary and secondary antibodies was carried out at room temperature for 45 min each. After washing, coverslips were mounted on slides and examined under a Zeiss LSM410 laser confocal microscope (Thornwood, NY).

FACS Analysis of N-cadherin Expression in Cocultures

Melanoma cells (3×10^5) were labeled with Cell Tracker Green (Molecular Probes) before coculture with an endothelial monolayer (3×10^5 cells) for 5 h. The cells were dissociated with 5 mM EDTA, incubated with N-cadherin mAb (GC4), followed by phycoerythrin-conjugated goat anti-mouse F(ab')₂ for 15 min each, and fixed in 3.5% paraformaldehyde. FACS analysis was performed to examine the level of N-cadherin on the cell surface of melanoma cells and endothelial cells.

Immunoprecipitation, Subcellular Fractionation, and Immunoblot Analysis

For biochemical analysis, 4×10^6 endothelial cells were cultured on a Matrigel-coated 60-mm plate in endothelial medium containing 10 ng/ml TNF- α for 12 h. The cells were washed three times and then incubated in fresh endothelial medium. An equal number of melanoma cells was deposited on the endothelial monolayer and cocultured for 5 h before lysis. To prepare samples for 0 h of coculture, melanoma cells and endothelial cells were cultured on Matrigel and collected separately. The cell lysates were mixed together before further analysis. Subcellular fractionation was performed to obtain membrane and cytosolic fractions or nuclear and postnuclear fractions according to published protocols (Dignam *et al.*, 1983; Thoreson *et al.*, 2000). For immunoprecipitation, samples were solubilized in lysis buffer (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, 1% NP-40, and a protease inhibitor cocktail). The supernatant was incubated overnight with 3 μ g of primary antibody at 4°C and then incubated with protein A-Sepharose beads (40 μ l of slurry made up of 10 mg dry beads) for 1 h at 4°C. The beads were washed three times in lysis buffer and boiled for 5 min before SDS-PAGE. For immunoblot analysis, protein samples were separated by SDS-PAGE and transferred to nitrocellulose membrane. Blots were probed with different antibodies and developed with ECL reagents. For quantification, the




Figure 1. Immunolocalization of N-cadherin during transendothelial migration of melanoma cells. WM239 melanoma cells were prelabeled with Cell Tracker Orange before seeding on a HMVEC monolayer. Cells were fixed at 1 h (a, e, and f), 3 h (c and g), or 5 h (b, d, and h) and then stained with mAb against VE-cadherin (a and b), E-cadherin (d), and N-cadherin (using mAb clone 32; c and e–h). Serial optical sections were taken by laser scanning confocal microscope, and selected sections were displayed with the distance of each section from the ventral surface shown in the right lower corner. The heterotypic contacts are indicated by arrows and the homotypic contacts between endothelial cells are indicated by arrowheads. The schematic drawing above or below each panel represents the stage of transendothelial migration shown in the micrograph. The sequential events are shown in e–h: (e) confocal image of a melanoma cell at the initial stage of attachment on the endothelium; (f) image of a melanoma cell beginning to send out membrane protrusions to invade the endothelial junction; (g) image of a melanoma cell with a large pseudopodium inserted into the endothelial junction; and (h) image of a melanoma cell intercalated between endothelial cells. Bars, 10 μm .

blots were analyzed in a Fluor-S imager (Bio-Rad, Hercules, CA) and quantification was based on the pixel value of each band after background subtraction.

Luciferase Reporter Assay

The TCF reporter plasmid kit, which contained the TOPflash, FOPflash, and TK vectors, was purchased from Upstate Biotechnology (Lake Placid, NY). Melanoma cells were cotransfected transiently with TOPflash (or FOPflash) and TK vectors at a ratio of 30:1 using Lipofectamine 2000. Two days later, melanoma cells (3×10^5) were seeded on top of a HMVEC monolayer (3×10^5 cells) in a Matrigel-coated 24-well plate. Cells were collected at different times and analyzed using the Promega's dual luciferase reporter assay system (Madison, WI). Luminescence was measured using a Berthold Lumat LB luminometer. The ratio between firefly luciferase activity (TOPflash or FOPflash) and Renilla luciferase activity (TK vector) was used to estimate changes in β -catenin-mediated transcription. The fold increase in luciferase activity was calculated by normalizing the data to that at 0 h of coculture.

RNA Isolation and Northern Blot Analysis

Melanoma cells (4×10^6 cells) were cultured on an endothelial monolayer (4×10^6 cells). Total RNA was isolated from 0- or 5-h cocultures using the RNeasy Mini Kit (Qiagen, Mississauga, Ontario, Canada). Equal amounts of total RNA ($\sim 3 \mu\text{g}$ per lane) were separated in 1.2% agarose gel containing 2% formaldehyde and then transferred onto positively charged nylon membranes. DIG-labeled RNA probes against CD44, N-cadherin, and MT1-MMP were generated using RT-PCR and in vitro transcription. RNA blots were

probed using the DIG-detection starter kit of Roche (Indianapolis, IN) according to the manufacturer's protocol.

RESULTS

N-cadherin Is Up-regulated and Concentrated in Heterotypic Contacts during Transendothelial Migration of Melanoma Cells

To determine whether N-cadherin was involved in transendothelial migration, labeled melanoma cells were deposited on a monolayer of endothelial cells. As previously reported (Voura *et al.*, 1998), melanoma cells attached on the endothelium and sent out membrane protrusions around their contact regions within 1 h. Pseudopodia were used to invade the endothelial junction and melanoma cells became intercalated between endothelial cells by 3 h. More than 50% of melanoma cells were spreading on the Matrigel by 5 h and they were scored as transmigrated cells.

The cocultures were fixed and stained with antibodies against different cadherins. VE-cadherin staining demarcated the endothelial junction, but it was distinctly absent from the heterotypic contact between endothelial cells and

melanoma cells (Figure 1a, b). N-cadherin was the major cadherin species in the heterotypic contact during transendothelial migration (Figure 1c). N-cadherin was diffusely distributed on the surface of endothelial cells and absent from the endothelial junction as reported by Salomon *et al.* (1992). E-cadherin showed only background level of staining and was not detectable in heterotypic contacts (Figure 1d).

Attachment of melanoma cells on the endothelium quickly induced redistribution of N-cadherin to the cell-cell contact regions (Figure 1e). Initial cell-cell interactions stimulated the melanoma cells to form membrane blebs and pseudopodia around the contact regions and N-cadherin staining was observed at the tips of these protrusions (Figure 1f). As the melanoma cell began its transmigration process by inserting pseudopodia into the endothelial junction, intense staining of N-cadherin was present around the pseudopodium (Figure 1g). Subsequently, melanoma cells became intercalated between endothelial cells and N-cadherin staining was observed along the leading edges of surrounding endothelial cells which began to spread on top of the melanoma cell (Figure 1h).

To investigate whether cell-cell interactions had any effect on the expression of N-cadherin during transendothelial migration, melanoma cells were cultured on an endothelial monolayer. Immunoblots showed a time-dependent increase in N-cadherin level, with an averaged 1.6-fold increase at 5 h (Figure 2A). Melanoma cells and endothelial cells were also cultured separately on Matrigel to determine whether Matrigel had any effect on N-cadherin synthesis. The results show that Matrigel did not induce N-cadherin expression in either cell type. To determine which cell type up-regulated N-cadherin expression during coculture, endothelial cells and dye-labeled melanoma cells were cocultured for 5 h. Cells were collected and then probed with an anti-N-cadherin antibody. FACS analysis was carried out to determine the cell surface level of N-cadherin for each cell population. A twofold increase was observed in melanoma cells, but little change was detected in endothelial cells (Figure 2B). The data suggest that cell-cell interactions during transendothelial migration stimulated the expression of N-cadherin in melanoma cells, but not in endothelial cells.

Knockdown of N-cadherin Expression in Melanoma Cells Inhibits Transendothelial Migration

The above results suggest that N-cadherin may play a role in transendothelial migration of melanoma cells. To assess the importance of N-cadherin in this process, the expression of N-cadherin in WM239 melanoma cells was repressed by stable transfection with an N-cadherin antisense construct (Figure 3A). Clones (A35, A36, and A2), which showed 90, 60, and 0% reduction in N-cadherin expression, respectively, were subjected to the transendothelial migration assay. Maximum inhibition was observed for A35 cells, with 60% inhibition at 3 h and 45% inhibition at 5 h of coculture, whereas A36 cells had an intermediate level of inhibition and A2 cells underwent transendothelial migration as efficiently as WM239 cells. The results show a dose-dependent relationship between N-cadherin level and transendothelial migration (Figure 3B). To block the residual amounts of N-cadherin expressed on A35 and A36 cells, melanoma cells were incubated with the function-blocking anti-N-cadherin mAb GC4 before deposition on the endothelial monolayer. Transendothelial migration was inhibited by another 10–15% in both A35 and A36 cells (Figure 3C). The data support a key role for N-cadherin in the transendothelial migration of melanoma cells.




Figure 2. Up-regulation of N-cadherin during transendothelial migration of melanoma cells. (A) Immunoblots showing N-cadherin expression during coculture: (a) samples taken at different times of transendothelial migration (TEM); (b) WM239 melanoma cells (MC) or endothelial cells (EC) cultured separately either on plastic or on Matrigel for 5 h. Blots were also probed with antibody against actin as a loading control. (B) FACS analysis of the mixed cell population collected from cocultures. WM239 cells were labeled with Cell Tracker Green (CTG) and N-cadherin was stained with phycoerythrin (PE)-conjugated antibody (red). (a) Endothelial cells and (b) melanoma cells were collected separately before coculture and then stained for N-cadherin; (c) A mixture of endothelial cells and melanoma cells collected from 5-h coculture. The increase of PE fluorescence represents an increase in N-cadherin. Quantification of the N-cadherin level before coculture (white) and at 5 h of coculture (shaded) was carried out for both cell types: (d) endothelial cells; (e) melanoma cells.

β -catenin Dissociates from N-cadherin and Translocates into the Nucleus of Transmigrating Melanoma Cells

Because actively transmigrating melanoma cells must continuously “break” and “remake” contacts with the endothelial surface, immunofluorescence studies were carried out to investigate changes in the N-cadherin adhesion complexes that might accompany transendothelial migration. When cocultures were stained for the catenins, both N-cadherin and β -catenin were found to concentrate in the heterotypic contact upon adherence of melanoma cells on the endothelium (Figure 4A, a and d). Surprisingly, when melanoma cells began to transmigrate across the endothelial junction, most of the heterotypic contacts became devoid of β -catenin (Figure 4Ae). In contrast, p120^{cas} remained clearly detectable in the heterotypic contact (Figure 4Ag). When homotypic contacts between melanoma cells were examined, both β -catenin and p120^{cas} were observed (Figure 4A, f and h). On the other hand, γ -catenin staining was absent from both




Figure 3. Effects of N-cadherin knockdown by antisense RNA on transendothelial migration of melanoma cells. (A) WM239 cells were transfected with an N-cadherin antisense construct. Equal numbers of stable transfectants were analyzed by immunoblot against N-cadherin. Clones A35 and A36 showed ~90 and 60% reduction in N-cadherin expression, respectively, whereas A2 showed little change (marked by asterisks). (B) WM239 cells (□) and clones A2 (●), A35 (○), and A36 (▲) were subjected to the transendothelial migration assay. (C) To neutralize the remaining N-cadherin, the transfectants were preincubated in the absence (shaded bar) or presence (white bar) of the N-cadherin function blocking mAb GC4 before the transendothelial migration assay, and the number of transmigrated cells was scored at 5 h. The parental WM239 cells were included as control. Data represent the mean \pm SD (n = 9).

heterotypic and homotypic contacts (Figure 4A, i and j), although it was expressed in WM239 cells. These results indicate that both N-cadherin and β -catenin were concentrated in the heterotypic contact during the initial stages of cell attachment, but signals accompanying the initiation of transendothelial migration somehow led to the dissociation of β -catenin from the N-cadherin adhesion complex.

Further examination of the micrographs showed that β -catenin became detectable in the nucleus of melanoma cells during transendothelial migration and displayed a punctate pattern of staining (Figure 4Ae). To provide biochemical evidence for β -catenin redistribution, immunoprecipitation studies were carried out (Figure 4B). The 5-h coculture was compared with the 0-h time point when the two cell types were simply mixed in order to provide the reference value before cell-cell interactions during transendothelial migration. Quantitative analysis showed a 50% reduction in β -catenin coprecipitated with N-cadherin at 5 h of coculture, although the total cellular level of β -catenin did not change over time. Subcellular fractionation studies showed a significant reduction of β -catenin associated with the membrane fraction at 5 h of coculture, which was accompanied by a corresponding increase in the cytosolic level of β -catenin (Figure 4C). An analysis of the nuclear fraction obtained from cocultures showed a 5–6-fold increase in the nuclear level of β -catenin (Figure 4D). As a control, melanoma cells and endothelial cells were cultured separately on Matrigel and the results showed that Matrigel did not induce the nuclear translocation of β -catenin. Together, these

results suggest the presence of an active mechanism involved in the translocation of β -catenin from the N-cadherin adhesion complex to the nucleus.

β -catenin Signaling Is Activated during Transendothelial Migration of Melanoma Cells

What then is the role of β -catenin in the nucleus during melanoma cell transmigration? It is well known that β -catenin interacts with the LEF/TCF family of transcription factors upon activation of Wnt signaling pathway. To determine whether β -catenin had a role in gene regulation during transendothelial migration, WM239 cells were transiently transfected with the TOPflash vector that contained wild-type TCF/LEF binding sites before a luciferase reporter gene. Changes in β -catenin-mediated transcription were examined during transendothelial migration. A time-dependent increase in luciferase activity was observed, probably reflecting a steady increase in the nuclear accumulation of β -catenin (Figure 5A). In contrast, luciferase activity remained at the background level in cells transfected with the FOPflash vector, which contained mutated TCF/LEF binding sites. After background subtraction, there was a sixfold increase in luciferase activity at 5 h of coculture, correlating well with the 5–6-fold increase in nuclear β -catenin (see Figure 4D).

Northern blot analysis was carried out using total RNA isolated from cocultures of WM239 cells and endothelial cells at 0 and 5 h. The RNA blots were probed for transcripts of N-cadherin, and the known β -catenin target genes CD44 and MT1-MMP. Quantitative analysis showed that N-cadherin had a 2.5-fold increase and CD44 displayed a threefold increase at 5 h of coculture. However, only a 1.2-fold increase was observed for MT1-MMP (Figure 5B). To confirm that the increase was due to β -catenin signaling in transmigrating melanoma cells, A21 cells which expressed a dominant-negative form of β -catenin (see next section) was used in coculture. As expected, there was little increase in the transcript level of the β -catenin target genes in these cocultures (Figure 5B).

To assess the effects of N-cadherin-mediated adhesion on β -catenin-dependent gene activation, N-cadherin present on the surface of TOPflash-transfected WM239 cells was blocked using the anti-N-cadherin mAb GC4. Luciferase activity was reduced by 50% at 5 h of coculture (Figure 5A). Furthermore, A2 and A35 cells were transfected with the TOPflash vector. A35 cells, which expressed only 10% of the normal level of N-cadherin, were unable to sequester β -catenin in the cell membrane. When luciferase activity was assayed, they showed a higher level of background activation that remained relatively constant during the 5-h period of coculture. On the other hand, the luciferase activity of A2 cells, which expressed the normal level of N-cadherin, was comparable to that of the parental WM239 cells (Figure 5C). Together, these results suggest a role for N-cadherin-mediated cell adhesion in the activation of the β -catenin signaling pathway.

Activation of β -catenin Signaling Facilitates Transendothelial Migration of Melanoma Cells

To provide direct evidence that β -catenin signaling was a key component of the transendothelial migration process, we transfected WM239 cells with either a dominant-negative form or a dominant-active form of β -catenin (DasGupta *et al.*, 2002). The N-terminal domain of β -catenin contains GSK phosphorylation sites that will target β -catenin to the ubiquitin-proteasome degradation pathway when phosphorylated. The central armadillo repeats of β -catenin are respon-




Figure 4. β -catenin dissociation from the heterotypic contact and translocation into the nucleus of melanoma cells during transendothelial migration. (A) Confocal images showing the localization of N-cadherin and catenins during transendothelial migration of WM239 cells (labeled red). Coverslips of 1- and 5-h cocultures were fixed and stained for either N-cadherin (green; a–c) or one of the catenins (green; d–j). For β -catenin staining (d–f), a mouse mAb against dephosphorylated β -catenin was used. Heterotypic contacts are indicated by arrows and β -catenin staining in the nucleus is marked by an arrowhead. The staining patterns of these proteins in homotypic contacts of WM239 cells (c, f, h, and j) were included for comparison. Bars, 10 μ m. (B) Analysis of proteins coimmunoprecipitated with N-cadherin. (a) Equal numbers of melanoma cells and endothelial cells were used in cocultures. Protein samples were precipitated with anti-N-cadherin pAb and blots were probed for N-cadherin, β -catenin, and p120^{ctn}. A reduction in N-cadherin-associated β -catenin at 5 h of coculture is observed (asterisk). (b) Quantification of the coimmunoprecipitation data. The pixel value of β -catenin or p120^{ctn} was compared with that of N-cadherin and the ratios were normalized to that of the 0 h sample. Data represent the mean \pm SD ($n = 3$); 0 h sample (shaded bar), 5 h sample (white bar); values of the 5-h samples are compared with their respective 0-h sample; Student's t test, * $p < 0.05$, ** $p > 0.1$. (C) Immunoblots of subcellular fractions showing redistribution of β -catenin during coculture. (a) Cells were collected from 0- and 5-h coculture. The nuclei were removed from cell homogenates to yield the postnuclear fraction (P), which was further separated into membrane (M) and cytosolic (C) fractions. All fractions were adjusted to the same volume and equal volumes were subjected to SDS-PAGE for immunoblot analyses. Reduction in the level of β -catenin in the membrane fraction (asterisk) was accompanied by an increase in the cytosolic fraction. (b) Quantification of subcellular fractions: M/P (shaded bar), C/P (white bar); the 5-h values of β -catenin are significantly different from their respective 0-h value; * $p < 0.05$. (D) Nuclear translocation of β -catenin during coculture. (a) Cell homogenates derived from 0- or 5-h cocultures were separated into the nuclear fraction (N) and postnuclear fraction (P). Protein blots were probed with antibodies against β -catenin and N-cadherin. (b) Endothelial cells and melanoma cells were also cultured separately on either plastic or Matrigel. Equal volumes of these samples were subjected to immunoblot analysis against β -catenin (arrow). (c) Quantitative analysis to determine the relative level of β -catenin in the nuclear fractions.

sible for binding with cadherin and the TCF/LEF transcription factors, whereas the C-terminal domain is the major transactivation domain. In this study, a β -catenin construct lacking both N- and C-terminal domains (Δ N+ Δ C) served as a dominant-negative form, whereas β -catenin lacking the N-terminal domain (Δ N) served as a dominant-active form (Figure 6A). The N-terminus of these mutant forms was fused to a TAP-tag, so that their expression in stably transfected cell lines could be probed with mAb against protein A (Figure 6B). The distribution of the

dominant-negative and dominant-active forms of β -catenin in transfectants was examined by double immunofluorescence staining with antiprotein-A mAb and anti- β -catenin polyclonal antibody. Although both mutant forms of β -catenin showed cytoplasmic and nuclear localization, a higher level was observed in the nucleus (Figure 6C). Interestingly, these two mutant forms of β -catenin did not colocalize with the endogenous β -catenin at the cell-cell contact regions, suggesting that they might not compete with the endogenous β -catenin for N-cadherin binding.




Figure 5. Activation of β -catenin-mediated transcription during transendothelial migration. (A) WM239 cells were transiently transfected with TOPflash or FOPflash vector. The transfectants were preincubated with or without the N-cadherin function blocking antibody GC4 and then deposited on a HMVEC monolayer. Cells were collected at different time intervals and assayed for luciferase activity: FOPflash (striped bar), TOPflash (gray bar), TOPflash + GC4 (white bar). Data represent the mean \pm SD ($n = 9$). (B) Transcriptional up-regulation of N-cadherin and β -catenin target genes during transendothelial migration of melanoma cells. Total RNA was isolated from 0- or 5-h cocultures and equal amounts ($\sim 3 \mu\text{g}$) of RNA were subjected to Northern blot analysis using DIG-labeled RNA probes of CD44, MT1-MMP and N-cadherin. The 18S rRNA was shown as the loading control. For comparison, RNA blots derived from cocultures of A21 cells, which expressed a dominant-negative form of β -catenin, were also probed. (C) The N-cadherin knockdown clones A2 (gray bar) and A35 (white bar) were transiently transfected with the TOPflash vector and then cultured on a HMVEC monolayer. Cells were collected at different times for luciferase assay. Data represent the mean \pm SD ($n = 9$).

Clones F13 (ΔN) and A21 ($\Delta\text{N}+\Delta\text{C}$) expressed high levels of their respective mutant β -catenin and were chosen for luciferase assays (Figure 6D) and transendothelial migration assays (Figure 6E). At 5 h of coculture, F13 cells showed 25 and 35% increase in transendothelial migration and β -catenin-mediated transcription, respectively. In contrast, A21 cells showed 50% reduction in β -catenin-induced luciferase activity and 40% reduction in transendothelial migration. For comparison, control transfectants with the TAP vector were assayed and they showed levels of luciferase activity and transmigration efficiency similar to those of WM239 cells. The data indicate a close correlation between the level of β -catenin-mediated transcription and the transmigration process.

Tyrosine Phosphorylation of N-cadherin Regulates the Binding of β -catenin during Transendothelial Migration

It remains to be determined what regulates the interactions between β -catenin and N-cadherin during transendothelial migration. Tyrosine phosphorylation has been implicated in the regulation of cadherin- β -catenin interaction (Gumbiner, 2000; Nelson and Nusse, 2004). When immunofluorescence studies were carried out to localize phospho-tyrosine, relatively weak signals were detected in the heterotypic contacts between transmigrating melanoma cells and endothelial cells (Figure 7A). To maximally visualize the phospho-tyrosine staining, cocultures were treated with the tyrosine phosphatase inhibitor pervanadate for 5 min before fixation. In the presence of pervanadate, the phospho-tyrosine staining became clearly visible in the heterotypic contact regions of transmigrating cells as well as the homotypic contact regions between endothelial cells. Notably, phospho-tyrosine staining was absent in the heterotypic contact before transmigration took place even in the presence of pervanadate (Figure 7A). As a control, WM239 cells alone were stained for phospho-tyrosine. Punctate staining was ob-

served in the cytoplasm but absent from the cell-cell contact regions (Figure 7B).

To further assess the role of protein phosphorylation during transendothelial migration, cocultures were carried out in the presence of different kinase inhibitors. The protein tyrosine kinase inhibitor genistein eliminated the phospho-tyrosine signal and inhibited the dissociation of β -catenin from the N-cadherin complex at the heterotypic contact (Figure 7C). When the number of transmigrated cells was scored at 5 h of coculture, addition of genistein and calphostin C (protein kinase C [PKC] inhibitor) resulted in 60 and 35% inhibition, respectively (Figure 7D). However, the protein kinase A inhibitor H8 had no effect.

To determine which protein was phosphorylated, the N-cadherin complex was isolated by immunoprecipitation and the protein blot was probed with anti-phospho-tyrosine mAb. A tyrosine-phosphorylated band of ~ 130 kDa, consistent with the size of N-cadherin, was detected at 5 h of coculture, but not at 0 h (Figure 8A). Although similar levels of N-cadherin were detected in the precipitates of the 0- and 5-h coculture samples, the amount of coprecipitated β -catenin decreased by $\sim 50\%$ at 5 h. To confirm the identity of the 130-kDa band, immunoprecipitation was performed using the anti-phospho-tyrosine mAb, and the blot was probed with mAb against N-cadherin and β -catenin. The antibody pulled down N-cadherin but not β -catenin at 5 h of coculture (Figure 8A), confirming that N-cadherin contained phosphorylated tyrosine but β -catenin did not.

To assess whether there were other tyrosine-phosphorylated proteins in the heterotypic contact, A35 cells, which expressed $<10\%$ of the cellular level of N-cadherin than the parental WM239 cells, were examined. When A35 cells were examined using the transendothelial migration assay, neither N-cadherin nor phospho-tyrosine staining was detected in the heterotypic contact (Figure 8B). The data indicate that N-cadherin was the major component that became tyrosine-




Figure 6. Effects of the dominant-negative and dominant-active forms of β -catenin on transendothelial migration of melanoma cells. (A) Schematic drawings of mutant β -catenins: the dominant-active form contained a deletion in the N-terminal domain (ΔN mutant), whereas the dominant-negative form contained both N- and C-terminal deletions ($\Delta N + \Delta C$ mutant). (B) Expression of mutant β -catenins in WM239 cells. Stable transfectants were subjected to immunoblot analysis using a mAb against protein A. Clones F13 (ΔN mutant) and A21 ($\Delta N + \Delta C$ mutant) were chosen for further analysis. (C) Immunolocalization of mutant β -catenins in transfected cells. Cells were fixed and double stained with anti-protein A mAb and anti- β -catenin pAb. Bars, 10 μ m. (D) Effects of mutant β -catenins on β -catenin-mediated gene transcription. F13 and A21 cells were transiently transfected with a TOPflash vector before coculture on a HMVEC monolayer. Cells were collected at 5 h and assayed for luciferase activity. (E) Effects of mutant β -catenins on transendothelial migration of the transfectants. The percentage of transmigrated cells was scored at 5 h. Data in D and E represent the mean \pm SD (n = 9). The asterisks indicate a statistically significant difference when compared with the control (p < 0.05).

phosphorylated in the heterotypic contact regions between melanoma cells and endothelial cells.

DISCUSSION

Extravasation of cancer cells is a complicated event that involves both adhesive interactions and chemotaxis (Voura *et al.*, 2001; Ramjeesingh *et al.*, 2003). In this report, we show that N-cadherin plays a key role in the transendothelial migration of melanoma cells. It is concentrated in the heterotypic contact between melanoma cells and endothelial cells. However, N-cadherin forms a novel type of adhesion complex that lacks β -catenin, which allows the activation of β -catenin signaling in transmigrating melanoma cells. Transendothelial migration of melanoma cells is impaired by blocking N-cadherin synthesis via antisense transfection or by inhibiting β -catenin signaling via the expression of a dominant-negative β -catenin in melanoma cells. It becomes apparent that both N-cadherin-mediated adhesion and β -catenin signaling are involved in transendothelial migration of melanoma cells.

Endothelial cells express two major cadherins, VE- and N-cadherin. Despite the relative abundance of N- and VE-cadherin, only VE-cadherin is concentrated in the endothelial junction. VE-cadherin is known to exclude N-cadherin

from the endothelial junction via the juxtamembrane region of its cytoplasmic domain (Navarro *et al.*, 1998). Our previous work show localized dissolution of VE-cadherin from the endothelial junction upon the attachment of melanoma cells (Sandig *et al.*, 1997b; Voura *et al.*, 2000). As VE-cadherin moves out, N-cadherin can now move in and becomes concentrated in the heterotypic contact during melanoma cell transmigration. N-cadherin is also enriched on filopodia, pseudopodia, and the migrating front of melanoma cells, consistent with the observation that cadherin is clustered in the tip of membrane processes where it acts as traction points for spreading and migration (Sandig *et al.*, 1997a; Adams *et al.*, 1998; Vasioukhin *et al.*, 2000).

In many tumors, down-regulation of E-cadherin is accompanied by up-regulation of N-cadherin (Islam *et al.*, 1996; Nieman *et al.*, 1999; Li *et al.*, 2001). This cadherin switch has been suggested to represent an invasive phenotype of tumors as an enhanced level of N-cadherin has been found to promote cell motility, invasion, and survival (Nieman *et al.*, 1999; Hazan *et al.*, 2000; Li *et al.*, 2001; Suyama *et al.*, 2002). Our data show that interactions between endothelial cells and melanoma cells stimulate a twofold increase in N-cadherin in the melanoma cells. Because transmigrating melanoma cells are surrounded on all sides by endothelial cells, the increase in N-cadherin may facilitate the formation of




Figure 7. Tyrosine phosphorylation in the heterotypic contact and β -catenin redistribution. (A) Tyrosine phosphorylation of proteins in the heterotypic contact during transendothelial migration of WM239 cells (labeled red). Cocultures were fixed at 1 or 5 h with or without the prior addition of 1 mM pervanadate (+PV or -PV) for 5 min and then stained for phospho-tyrosine (green). Arrows indicate heterotypic contacts between melanoma cells and endothelial cells. (B) Lack of phospho-tyrosine staining in homotypic contacts of melanoma cells. WM239 cells were cultured on Matrigel, treated with 1 mM pervanadate for 5 min before fixation, and then double stained for β -catenin (red) and phospho-tyrosine (P-Tyr; green). (C) Inhibition of β -catenin dissociation from the heterotypic contact by genistein. WM239 cells (red) were cultured on top of a HMVEC monolayer in the presence of genistein (10 μ M). Coverslips were fixed and stained with mAb against phospho-tyrosine, N-cadherin, or β -catenin. Arrows indicate heterotypic contacts which showed N-cadherin (green) and β -catenin (green) localization but lacked phospho-tyrosine staining. Similar experiments were carried out with calphostin C and H8, but they had no effect (unpublished data). (D) Effects of protein kinase inhibitors on melanoma cell transmigration. Transendothelial migration assays were carried out in the presence of genistein, calphostin C or H8, and transmigrated cells were scored at 5 h. Data represent the mean \pm SD ($n = 9$). Bars, 10 μ m.

heterotypic contacts with the endothelial cells. N-cadherin is apparently up-regulated at the transcription level by β -catenin signaling in transmigrating melanoma cells. In contrast, the level of N-cadherin in the endothelial cells remains relatively constant. This may be due to the lack of β -catenin signaling in endothelial cells because only a small area of the endothelial membrane is engaged in heterotypic contact formation and, therefore, most of the β -catenin would remain sequestered in the homotypic endothelial junction via interaction with VE-cadherin.


It is well known that β -catenin maintains the stability of adhesion complexes mediated by classical cadherins by linking them to the actin cytoskeleton. Mutations in β -catenin or α -catenin that disrupt this linkage eliminate adhesiveness mediated by cadherin (Kawanishi *et al.*, 1995; Bullions *et al.*, 1997). Also, cadherin loses its binding activity after truncation or deletion of its cytoplasmic domain (Nagafuchi and Takeichi, 1988; Ozawa *et al.*, 1990). In our studies, coimmunolocalization and coimmunoprecipitation studies have been used to confirm the association of β -catenin with N-cadherin at the initial stage of cell attachment. However, in

subsequent stages of transmigration, the level of β -catenin associated with the heterotypic contact or the N-cadherin complex is reduced significantly. Cadherins without linkage to the actin cytoskeleton may undergo weaker interactions. Because tight adhesion is believed to be detrimental to cell migration (Nakagawa and Takeichi, 1998), weaker adhesive interactions may promote the transmigration of melanoma cell.

In the N-cadherin immunoprecipitates, a substantial amount of p120^{ctn} was also present. p120^{ctn} binds to the juxtamembrane region of cadherins and has been shown to facilitate the clustering of C-cadherin even in the absence of β -catenin binding (Yap *et al.*, 1998) and enhance E-cadherin-mediated adhesion (Thoreson *et al.*, 2000). It is possible that p120^{ctn} may contribute to the stability of the N-cadherin complex and prevent its dissolution in the heterotypic contact during melanoma cell transmigration.

In contrast to reports showing that elevated levels of either N- or E-cadherin can inhibit Wnt signaling by sequestering β -catenin (Sadot *et al.*, 1998; Nelson and Nusse, 2004), we have found that β -catenin dissociates from the N-cad-

Figure 8. Phosphorylation of N-cadherin in the heterotypic contact during transendothelial migration of melanoma cells. (A) Tyrosine phosphorylation of N-cadherin during transendothelial migration of WM239 cells. Equal number of melanoma cells and endothelial cells were cocultured for 5 h. Cells were collected and subjected to coimmunoprecipitation using either anti-N-cadherin pAb or anti-phospho-tyrosine mAb. The immunoprecipitates were probed with mAb against N-cadherin, β -catenin, or phospho-tyrosine. The arrows indicate N-cadherin (~ 130 kDa) and the arrowheads indicate β -catenin (~ 92 kDa). (B) Loss of phospho-tyrosine staining in heterotypic contacts during transendothelial migration of N-cadherin knockdown melanoma cells (labeled with asterisks). A35 cells were cultured on a HMVEC monolayer for 5 h. Coverslips were fixed and stained with mAb against N-cadherin or phospho-tyrosine, with or without pervanadate pretreatment (+PV or -PV). Arrows indicate the absence of N-cadherin or phospho-tyrosine staining in the heterotypic contact and the arrowheads indicate phospho-tyrosine staining in the endothelial junction. Bars, 10 μ m.



herin adhesion complex and activates the β -catenin signaling pathway in transmigrating melanoma cells despite an up-regulation of N-cadherin. The β -catenin signaling pathway has been implicated in cancer metastasis (Smalley and Dale, 1999; Peifer and Polakis, 2000). More than 80% of human colorectal carcinomas contain mutations in either APC or β -catenin, leading to constitutive activation of the β -catenin signaling pathway (Bienz and Clevers, 2000). Nuclear β -catenin is frequently observed in colon cancer cells at the invasion front (Brabletz *et al.*, 2001; Hiendlmeyer *et al.*, 2004), suggesting the possible involvement of β -catenin signaling in invasion. It is of interest to note that among the known β -catenin target genes are those that code for proteins involved in the invasive process of cancer cells, such as CD44 (Wielenga *et al.*, 1999), uPA (Hiendlmeyer *et al.*, 2004), uPAR (Mann *et al.*, 1999), MMP-7 (Brabletz *et al.*, 1999), and MT1-MMP (Takahashi *et al.*, 2002), and we have confirmed the up-regulation of CD44. Up-regulation of invasion-related genes should augment the transmigration process by promoting invasion of the basement membrane. Indeed, blocking β -catenin signaling by expressing a dominant-negative β -catenin results in significant inhibition of transendothelial migration, whereas the expression of a dominant-active β -catenin promotes transmigration.

Normally, excess β -catenin will be targeted for degradation in the absence of the Wnt signal (Jamora and Fuchs, 2002). However, the level of β -catenin does not change in our coculture system, suggesting that the melanoma cells might utilize a different mechanism to stabilize β -catenin. In addition, both dominant-active and dominant-negative mutants of β -catenin show preferential nuclear localization. These two mutant forms lack the N-terminal degradation signal, thus allowing them to escape degradation by the ubiquitin-proteasome pathway, whereas their armadillo repeats might facilitate their nuclear translocation (Fagotto *et al.*, 1998; Yokoya *et al.*, 1999). Sequestration of the mutant β -catenins in the nucleus would lower their cytoplasmic level, rendering them unable to compete with the endogenous β -catenin for binding to the membrane-associated N-

cadherin. However, their high concentration in the nucleus should allow them to compete efficiently with the endogenous β -catenin for binding to the target genes.

The dissociation of β -catenin is a postattachment event, concomitant with the transmigration process. We speculate that heterotypic cell-cell interactions generate a signal that is capable of triggering a cascade of events, which in turn leads to β -catenin dissociation. Although the nature of this signal is not known, our results demonstrate a correlation between tyrosine phosphorylation of N-cadherin and the dissociation of β -catenin from the adhesion complex. In contrast to E-, N-, or P-cadherins which are either not phosphorylated or poorly phosphorylated, VE-cadherin can be heavily phosphorylated on its tyrosine residues under physiological conditions (Lampugnani *et al.*, 1997). PECAM-1 is also known to contain phospho-tyrosine (Osawa *et al.*, 2002). However, both VE-cadherin and PECAM-1 are absent from heterotypic contacts (Sandig *et al.*, 1997b; Voura *et al.*, 2000), suggesting that the phospho-tyrosine staining in the heterotypic contact is not due to either of these adhesion proteins. Notably, the phospho-tyrosine staining disappears from the heterotypic contacts when N-cadherin is knocked down in melanoma cells, indicating that the phospho-tyrosine residues are associated with N-cadherin.

It is likely that heterotypic cell-cell interactions between melanoma cells and endothelial cells stimulate the activity of a protein tyrosine kinase, which phosphorylates the cytoplasmic domain of N-cadherin, thereby destabilizing the binding of β -catenin to N-cadherin. In coculture studies, addition of genistein inhibits tyrosine phosphorylation of N-cadherin and prevents β -catenin dissociation. Genistein also inhibits transendothelial migration by 60%.

Although tyrosine phosphorylation of N-cadherin is observed, we have failed to detect tyrosine-phosphorylated β -catenin. Our results, therefore, stand in striking contrast to what is commonly held regarding the regulation of cadherin- β catenin interaction. Tyrosine phosphorylation of β -catenin has long been believed to perturb cadherin-mediated adhesion by disrupting cadherin- β -catenin interaction.

β -catenin becomes phosphorylated, resulting in the disruption of cell-cell adhesion, when cells are transfected with v-Src, treated with tyrosine phosphatase inhibitor or stimulated with growth factors (Volberg *et al.*, 1992; Behrens *et al.*, 1993; Shibamoto *et al.*, 1994; Kinch *et al.*, 1995; Balsamo *et al.*, 1996; Nelson and Nusse, 2004). Tyrosine-phosphorylated β -catenin shows reduced binding to E-cadherin *in vitro*, and crystallography studies suggest that tyrosine phosphorylation of β -catenin may disrupt its interaction with E-cadherin through steric effects (Roura *et al.*, 1999; Huber and Weis, 2001). However, little is known about the effect of tyrosine phosphorylation of N-cadherin on β -catenin binding because N-cadherin is poorly phosphorylated on tyrosine residues.

Recently, an active c-Src mutant has been reported to induce tyrosine phosphorylation of cadherin but not β -catenin (Irby and Yeatman, 2002). N-cadherin possesses several tyrosine residues that are potential substrates of Src and several of them reside within the β -catenin binding region. It would be of interest to determine whether Src activation is responsible for the tyrosine phosphorylation of N-cadherin, leading to the subsequent dissociation of β -catenin during melanoma cell transmigration.

Using other kinase inhibitors, we have found that H8 does not inhibit transendothelial migration, suggesting that protein kinase A may not play a role here. On the other hand, PKC has been implicated in cancer cell chemotaxis, migration, and invasion (Masur *et al.*, 2001; Koivunen *et al.*, 2004; Sun *et al.*, 2005). Indeed, calphostin C inhibits transendothelial migration by almost 40%, suggesting that multiple signaling pathways are activated during extravasation. Future studies involving the identification of the kinases and phosphatases that regulate the phosphorylation state of N-cadherin and other signaling components should help define the signaling pathways involved in cancer cell transendothelial migration.

ACKNOWLEDGMENTS

We are grateful to Drs. Rama Khoka and Jaro Sodek for invaluable advice and discussion. We thank Dr. Bertrand Seraphin (CNRS, France) for the original TAP-tag construct, Dr. Keith Johnson (University of Toledo, Toledo, OH) for the pPUR construct, Dr. Meenhard Herlyn (Wistar Institute, Philadelphia, PA) for the melanoma cell lines WM239 and WM35, and Dr. Bernard Schimmer (University of Toronto) for assistance in the luciferase assay. This work was supported by an operating grant from the Canadian Institutes of Health Research (MT-14703). J.Q. is supported in part by a Connaught Entrance Scholarship and a University of Toronto Open Scholarship. N.C. was supported by a Canadian Institute of Health Research Studentship. J.W. was the recipient of a scholarship from the Shandong Research Council of PR China.

REFERENCES

Adams, C. L., Chen, Y. T., Smith, S. J., and Nelson, W. J. (1998). Mechanisms of epithelial cell-cell adhesion and cell compaction revealed by high-resolution tracking of E-cadherin-green fluorescent protein. *J. Cell Biol.* 142, 1105–1119.

Aigner, S., Ramos, C. L., Hafezi-Moghadam, A., Lawrence, M. B., Friederichs, J., Altevogt, P., and Ley, K. (1998). CD24 mediates rolling of breast carcinoma cells on P-selectin. *FASEB J.* 12, 1241–1251.

Anastasiadis, P. Z., and Reynolds, A. B. (2000). The p120 catenin family: complex roles in adhesion, signaling and cancer. *J. Cell Sci.* 113, 1319–1334.

Balsamo, J., Leung, T., Ernst, H., Zanin, M. K., Hoffman, S., and Lilien, J. (1996). Regulated binding of PTP1B-like phosphatase to N-cadherin: control of cadherin-mediated adhesion by dephosphorylation of β -catenin. *J. Cell Biol.* 134, 801–813.

Behrens, J., Vakaet, L., Friis, R., Winterhager, E., Van Roy, F., Mareel, M. M., and Birchmeier, W. (1993). Loss of epithelial differentiation and gain of invasiveness correlates with tyrosine phosphorylation of the E-cadherin/ β -catenin complex in cells transformed with a temperature-sensitive v-SRC gene. *J. Cell Biol.* 120, 757–766.

Behrens, J., von Kries, J. P., Kuhl, M., Bruhn, L., Wedlich, D., Grosschedl, R., and Birchmeier, W. (1996). Functional interaction of β -catenin with the transcription factor LEF-1. *Nature* 382, 638–642.

Bienz, M., and Clevers, H. (2000). Linking colorectal cancer to Wnt signaling. *Cell* 103, 311–320.

Brabletz, T., Jung, A., Dag, S., Hlubek, F., and Kirchner, T. (1999). β -catenin regulates the expression of the matrix metalloproteinase-7 in human colorectal cancer. *Am. J. Pathol.* 155, 1033–1038.

Brabletz, T., Jung, A., Reu, S., Porzner, M., Hlubek, F., Kunz-Schughart, L. A., Knuechel, R., and Kirchner, T. (2001). Variable β -catenin expression in colorectal cancers indicates tumor progression driven by the tumor environment. *Proc. Natl. Acad. Sci. USA* 98, 10356–10361.

Brandt, B. H. *et al.* (1999). c-erbB-2/EGFR as dominant heterodimerization partners determine a motogenic phenotype in human breast cancer cells. *FASEB J.* 13, 1939–1949.

Brenner, W., Langer, P., Oesch, F., Edgell, C. J., and Wieser, R. J. (1995). Tumor cell-endothelium adhesion in an artificial venule. *Anal. Biochem.* 225, 213–219.

Bullions, L. C., Nottelman, D. A., Chung, L. S., and Levine, A. J. (1997). Expression of wild-type α -catenin protein in cells with a mutant α -catenin gene restores both growth regulation and tumor suppressor activities. *Mol. Cell. Biol.* 17, 4501–4508.

Chambers, A. F., Schimdt, E. E., MacDonald, I. C., Morris, V. L., and Groom, A. C. (1992). Early steps in hematogenous metastasis of B16F1 melanoma cells in chick embryos studied by high resolution intravital videomicroscopy. *J. Natl. Cancer Inst.* 84, 797–803.

Christofori, G. (2003). Changing neighbours, changing behaviour: cell adhesion molecule-mediated signalling during tumour progression. *EMBO J.* 22, 2318–2323.

DasGupta, R., Rhee, H., and Fuchs, E. (2002). A developmental conundrum: a stabilized form of β -catenin lacking the transcriptional activation domain triggers features of hair cell fate in epidermal cells and epidermal cell fate in hair follicle cells. *J. Cell Biol.* 158, 331–344.

Dignam, J. D., Lebovitz, R. M., and Roeder, R. G. (1983). Accurate transcription initiation by RNA polymerase II in a soluble extract from isolated mammalian nuclei. *Nucleic Acids Res.* 11, 1475–1489.

Fagotto, F., Gluck, U., and Gumbiner, B. M. (1998). Nuclear localization signal-independent and importin/karyopherin-independent nuclear import of β -catenin. *Curr. Biol.* 8, 181–190.

Fidler, I. J. (2003). The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat. Rev. Cancer* 3, 453–458.

Giavazzi, R., Foppolo, M., Dossi, R., and Remuzzi, A. (1993). Rolling and adhesion of human tumor cells on vascular endothelium under physiological flow conditions. *J. Clin. Invest.* 92, 3038–3044.

Gumbiner, B. M. (2000). Regulation of cadherin adhesive activity. *J. Cell Biol.* 148, 399–404.

Hazan, R. B., Phillips, G. R., Qiao, R. F., Norton, L., and Aaronson, S. A. (2000). Exogenous expression of N-cadherin in breast cancer cells induces cell migration, invasion, and metastasis. *J. Cell Biol.* 148, 779–790.

Hiendlmeyer, E. *et al.* (2004). β -catenin up-regulates the expression of the urokinase plasminogen activator in human colorectal tumors. *Cancer Res.* 64, 1209–1214.

Huber, A. H., and Weis, W. I. (2001). The structure of the β -catenin/E-cadherin complex and the molecular basis of diverse ligand recognition by β -catenin. *Cell* 105, 391–402.

Ilan, N., and Madri, J. A. (2003). PECAM-1, old friend, new partners. *Curr. Opin. Cell Biol.* 15, 515–524.

Irby, R. B., and Yeatman, T. J. (2002). Increased Src activity disrupts cadherin/catenin-mediated homotypic adhesion in human colon cancer and transformed rodent cells. *Cancer Res.* 62, 2669–2674.

Islam, S., Carey, T. E., Wolf, G. T., Wheelock, M. J., and Johnson, K. R. (1996). Expression of N-cadherin by human squamous carcinoma cells induces a scattered fibroblastic phenotype with disrupted cell-cell adhesion. *J. Cell Biol.* 135, 1643–1654.

Jamora, C., and Fuchs, E. (2002). Intercellular adhesion, signalling and the cytoskeleton. *Nat. Cell Biol.* 4, E101–E108.

Kawanishi, J., Kato, J., Sasaki, K., Fujii, S., Watanabe, N., and Niitsu, Y. (1995). Loss of E-cadherin-dependent cell-cell adhesion due to mutation of the β -catenin gene in a human cancer cell line, HSC-39. *Mol. Cell. Biol.* 15, 1175–1181.

Kinch, M. S., Clark, G. J., Der, C. J., and Burridge, K. (1995). Tyrosine phosphorylation regulates the adhesions of ras-transformed breast epithelia. *J. Cell Biol.* 130, 461–471.

- Koivunen, J., Aaltonen, V., Koskela, S., Lehenkari, P., Laato, M., and Peltonen, J. (2004). PKC α/β inhibitor Go6976 promotes formation of cell junctions and inhibits invasion of urinary bladder carcinoma cells. *Cancer Res.* 64, 5693–5701.
- Lampugnani, M. G., Corada, M., Andriopoulou, P., Esser, S., Risau, W., and Dejana, E. (1997). Cell confluence regulates tyrosine phosphorylation of adherens junction components in endothelial cells. *J. Cell Sci.* 110, 2065–2077.
- Li, G., and Herlyn, M. (2000). Dynamics of intercellular communication during melanoma development. *Mol. Med. Today* 6, 163–169.
- Li, G., Satyamoorthy, K., and Herlyn, M. (2001). N-cadherin-mediated intercellular interactions promote survival and migration of melanoma cells. *Cancer Res.* 61, 3819–3825.
- Luscinskas, F. W., Ding, H., and Lichtman, A. H. (1995). P-selectin and vascular cell adhesion molecule 1 mediate rolling and arrest, respectively, of CD4+ T lymphocyte on tumor necrosis factor-activated vascular endothelium under flow. *J. Exp. Med.* 181, 1179–1186.
- Mann, B. *et al.* (1999). Target genes of β -catenin-T cell-factor/lymphoid-enhancer-factor signaling in human colorectal carcinomas. *Proc. Natl. Acad. Sci. USA* 96, 1603–1608.
- Masur, K., Lang, K., Niggemann, B., Zanker, K. S., and Entschladen, F. (2001). High PKC α and low E-cadherin expression contribute to high migratory activity of colon carcinoma cells. *Mol. Biol. Cell* 12, 1973–1982.
- Nagafuchi, A., and Takeichi, M. (1988). Cell binding function of E-cadherin is regulated by the cytoplasmic domain. *EMBO J.* 7, 3679–3684.
- Nagafuchi, A. (2001). Molecular architecture of adherens junctions. *Curr. Opin. Cell Biol.* 13, 600–603.
- Nakagawa, S., and Takeichi, M. (1998). Neural crest emigration from the neural tube depends on regulated cadherin expression. *Development* 125, 2963–2971.
- Navarro, P., Ruco, L., and Dejana, E. (1998). Differential localization of VE- and N-cadherins in human endothelial cells: VE-cadherin competes with N-cadherin for junctional localization. *J. Cell Biol.* 140, 1475–1484.
- Nelson, W. J., and Nusse, R. (2004). Convergence of Wnt, β -catenin, and cadherin pathways. *Science* 303, 1483–1487.
- Nieman, M. T., Prudoff, R. S., Johnson, K. R., and Wheelock, M. J. (1999). N-cadherin promotes motility in human breast cancer cells regardless of their E-cadherin expression. *J. Cell Biol.* 147, 631–644.
- Orr, F. W., Wang, H. H., Lafrenie, R. M., Scherbarth, S., and Nance, D. M. (2000). Interactions between cancer cells and the endothelium in metastasis. *J. Pathol.* 190, 310–329.
- Osawa, M., Masuda, M., Kusano, K., and Fujiwara, K. (2002). Evidence for a role of platelet endothelial cell adhesion molecule-1 in endothelial cell mechanosignal transduction: is it a mechanoresponsive molecule? *J. Cell Biol.* 158, 773–785.
- Ozawa, M., Ringwald, M., and Kemler, R. (1990). Uvomorulin-catenin complex formation is regulated by a specific domain in the cytoplasmic region of the cell adhesion molecule. *Proc. Natl. Acad. Sci. USA* 87, 4246–4250.
- Pantel, K., and Brakenhoff, R. H. (2004). Dissecting the metastatic cascade. *Nat. Rev. Cancer* 4, 448–456.
- Peifer, M., and Polakis, P. (2000). Wnt signaling in oncogenesis and embryogenesis—a look outside the nucleus. *Science* 287, 1606–1609.
- Ramjeesingh, R., Leung, R., and Siu, C. H. (2003). Interleukin-8 secreted by endothelial cells induces chemotaxis of melanoma cells through the chemokine receptor CXCR1. *FASEB J.* 17, 1292–1304.
- Rigaut, G., Shevchenko, A., Rutz, B., Wilm, M., Mann, M., and Seraphin, B. (1999). A generic protein purification method for protein complex characterization and proteome exploration. *Nat. Biotechnol.* 17, 1030–1032.
- Roura, S., Miravet, S., Piedra, J., Garcia de Herreros, A., and Dunach, M. (1999). Regulation of E-cadherin/catenin association by tyrosine phosphorylation. *J. Biol. Chem.* 274, 36734–36740.
- Sadot, E., Simcha, I., Shtutman, M., Ben-Ze'ev, A., and Geiger, B. (1998). Inhibition of β -catenin-mediated transactivation by cadherin derivatives. *Proc. Natl. Acad. Sci. USA* 95, 15339–15344.
- Salomon, D., Ayalon, O., Patel-King, R., Hynes, R. O., and Geiger, B. (1992). Extrajunctional distribution of N-cadherin in cultured human endothelial cells. *J. Cell Sci.* 102, 7–17.
- Sandig, M., Kalnins, V. I., and Siu, C. H. (1997a). Role of NCAM, cadherins, and microfilaments in cell-cell contact formation in TM4 immature mouse sertoli cells. *Cell Motil. Cytoskelet.* 36, 149–163.
- Sandig, M., Voura, E. B., Kalnins, V. I., and Siu, C. H. (1997b). Role of cadherins in the transendothelial migration of melanoma cells in culture. *Cell Motil. Cytoskelet.* 38, 351–364.
- Shibamoto, S., Hayakawa, M., Takeuchi, K., Hori, T., Oku, N., Miyazawa, K., Kitamura, N., Takeichi, M., and Ito, F. (1994). Tyrosine phosphorylation of β -catenin and plakoglobin enhanced by hepatocyte growth factor and epidermal growth factor (EGF) in human carcinoma cells. *Cell Adhes. Commun.* 1, 295–305.
- Smalley, M. J., and Dale, T. C. (1999). Wnt signaling in mammalian development and cancer. *Cancer Metastasis Rev.* 18, 215–230.
- Sun, R., Gao, P., Chen, L., Ma, D., Wang, J., Oppenheim, J. J., and Zhang, N. (2005). PKC ζ is required for EGF-induced chemotaxis of human breast cancer cells. *Cancer Res.* 65, 1433–1441.
- Suyama, K., Shapiro, I., Guttman, M., and Hazan, R. B. (2002). A signaling pathway leading to metastasis is controlled by N-cadherin and the FGF receptor. *Cancer Cell* 2, 301–314.
- Takahashi, M., Tsunoda, T., Seiki, M., Nakamura, Y., and Furukawa, Y. (2002). Identification of membrane-type matrix metalloproteinase-1 as a target of the β -catenin/Tcf4 complex in human colorectal cancers. *Oncogene* 21, 5861–5867.
- Takeichi, M. (1995). Morphogenetic roles of classic cadherins. *Curr. Opin. Cell Biol.* 7, 619–627.
- Thoreson, M. A., Anastasiadis, P. Z., Daniel, J. M., Ireton, R. C., Wheelock, M. J., Johnson, K. R., Hummingbird, D. K., and Reynolds, A. B. (2000). Selective uncoupling of p120^{cas} from E-cadherin disrupts strong adhesion. *J. Cell Biol.* 148, 189–202.
- Tomita, K., van Bokhoven, A., van Leenders, G. J., Ruijter, E. T., Jansen, C. F., Bussemakers, M. J., and Schalken, J. A. (2000). Cadherin switching in human prostate cancer progression. *Cancer Res.* 60, 3650–3654.
- Vasioukhin, V., Bauer, C., Yin, M., and Fuchs, E. (2000). Directed actin polymerization is the driving force for epithelial cell-cell adhesion. *Cell* 100, 209–219.
- Vestweber, D. (2002). Regulation of endothelial cell contacts during leukocyte extravasation. *Curr. Opin. Cell Biol.* 14, 587–593.
- Volberg, T., Zick, Y., Dror, R., Sabanay, I., Gilon, C., Levitzki, A., and Geiger, B. (1992). The effect of tyrosine-specific protein phosphorylation on the assembly of adherens-type junctions. *EMBO J.* 11, 1733–1742.
- Voura, E. B., Sandig, M., Kalnins, V. I., and Siu, C. H. (1998). Cell shape changes and cytoskeleton reorganization during transendothelial migration of human melanoma cells. *Cell Tissue Res.* 293, 375–387.
- Voura, E. B., Chen, N., and Siu, C. H. (2000). Platelet-endothelial cell adhesion molecule-1 (CD31) redistributes from the endothelial junction and is not required for the transendothelial migration of melanoma cells. *Clin. Exp. Metastasis* 18, 527–532.
- Voura, E. B., Ramjeesingh, R. A., Montgomery, A. M., and Siu, C. H. (2001). Involvement of integrin $\alpha v \beta 3$ and cell adhesion molecule L1 in transendothelial migration of melanoma cells. *Mol. Biol. Cell* 12, 2699–2710.
- Wielenga, V. J., Smits, R., Korinek, V., Smit, L., Kielman, M., Fodde, R., Clevers, H., and Pals, S. T. (1999). Expression of CD44 in Apc and Tcf mutant mice implies regulation by the WNT pathway. *Am. J. Pathol.* 154, 515–523.
- Yap, A. S., Niessen, C. M., and Gumbiner, B. M. (1998). The juxtamembrane region of the cadherin cytoplasmic tail supports lateral clustering, adhesive strengthening, and interaction with p120^{cas}. *J. Cell Biol.* 141, 779–789.
- Yokoya, F., Imamoto, N., Tachibana, T., and Yoneda, Y. (1999). β -catenin can be transported into the nucleus in a Ran-unassisted manner. *Mol. Biol. Cell* 10, 1119–1131.